

# Algebraically positive matrices\*

Steve Kirkland<sup>a</sup>, Pu Qiao<sup>b</sup>, Xingzhi Zhan<sup>b†</sup>

<sup>a</sup>Department of Mathematics, University of Manitoba, Winnipeg, MB, Canada

<sup>b</sup>Department of Mathematics, East China Normal University, Shanghai 200241, China

## Abstract

We introduce the concept of algebraically positive matrices and investigate some basic properties, including a characterization, the index of algebraic positivity, and sign patterns that allow or require this property. We also pose two open problems.

**Key words.** Algebraically positive matrix, primitive matrix, irreducible matrix, sign pattern

**Mathematics subject classifications.** 15B48, 15B35

## 1 Introduction

A *positive (nonnegative) matrix* is a matrix all of whose entries are positive (nonnegative) real numbers. The notation  $A > 0$  means that  $A$  is a positive matrix. We introduce the following concept and study its basic properties.

**Definition.** A square real matrix  $A$  is said to be *algebraically positive* if there exists a real polynomial  $f$  such that  $f(A)$  is a positive matrix.

The motivation for this concept is twofold. First, we would like to extend the concept of primitive nonnegative matrices to algebraically positive matrices for general real matrices;

---

\*E-mail addresses: [Stephen.Kirkland@umanitoba.ca](mailto:Stephen.Kirkland@umanitoba.ca)(S.Kirkland), [235711gm@sina.com](mailto:235711gm@sina.com)(P.Qiao), [zhan@math.ecnu.edu.cn](mailto:zhan@math.ecnu.edu.cn)(X.Zhan). The research presented in this paper was supported in part by NSERC, the Shanghai SF grant 15ZR1411500 and the NSFC grant 11371145.

<sup>†</sup>Corresponding author.

second, we wish to let the negative entries and positive entries in a real matrix have equal status. Also, the action of polynomials on matrices is very common in matrix analysis ([1, Chapter V], [4, Chapter 6]) and even in graph theory [2, Section 5.3]. All the matrices in this paper are square unless otherwise specified, so we will omit the word “square”.

Recall that if an irreducible nonnegative matrix  $A$  has exactly  $k$  eigenvalues of modulus equal to the spectral radius of  $A$ , then  $k$  is called the *index of imprimitivity* of  $A$ . If  $k = 1$ ,  $A$  is said to be *primitive*; otherwise,  $A$  is *imprimitive*. It is known (see, e.g., [7, p.134]) that a nonnegative matrix  $A$  of order at least 2 is primitive if and only if there exists a positive integer  $m$  such that  $A^m$  is a positive matrix. Thus a primitive matrix is algebraically positive.

There is another related concept. A real matrix  $A$  is said to be *eventually positive* if there exists a positive integer  $p$  such that  $A^k$  is a positive matrix for all integers  $k \geq p$ . See [6] and the references therein for recent development and the history of this topic. Clearly, every eventually positive matrix is algebraically positive. Note that eventually positive matrices are a proper subclass of algebraically positive matrices. For example, the matrix  $A = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$  is algebraically positive since  $I - A > 0$ , but  $A$  is not eventually positive, since it is idempotent and has some negative entries.

Obviously, if a real matrix  $A$  is algebraically positive, then so are  $-A$  and  $P^T A P$  for any permutation matrix  $P$ .

We remark that the possible corresponding concepts of *algebraically nonnegative matrices* and *algebraically positive definite matrices* are not so compelling, since every real matrix is algebraically nonnegative by the Cayley-Hamilton theorem (see, e.g., [7, p.24]) while it is straightforward to see that any real symmetric matrix is algebraically positive definite.

We will study some basic properties of algebraically positive matrices, including a characterization of algebraic positivity, the index of algebraic positivity, and sign patterns. At the end we pose two open problems.

## 2 Main results

Let  $A$  be a real matrix of order  $n$  with characteristic polynomial  $h(x)$  and let  $f(x)$  be any real polynomial. By the division algorithm there exist real polynomials  $q(x)$  and  $r(x)$

with  $\deg r(x) \leq n - 1$  such that  $f(x) = h(x)q(x) + r(x)$ . The Cayley-Hamilton theorem asserts that  $h(A) = 0$ . Thus  $f(A) = r(A)$ . It follows that to check whether a given real matrix of order  $n$  is algebraically positive we need only consider polynomials of degree less than or equal to  $n - 1$ .

Throughout we denote by  $I$  the identity matrix whose order will be clear from the context and by  $A^T$  the transpose of a matrix  $A$ . We write the elements of  $\mathbb{R}^n$  as column vectors. A *simple eigenvalue* is an eigenvalue of algebraic multiplicity 1.

**Theorem 1** *A real matrix is algebraically positive if and only if it has a simple real eigenvalue and corresponding left and right positive eigenvectors.*

**Proof.** Let  $A$  be an algebraically positive matrix and let  $f$  be a real polynomial such that  $f(A) > 0$ . Let  $\lambda_1, \dots, \lambda_n$  be the eigenvalues of  $A$ . The spectral mapping theorem [7, p.8] asserts that the eigenvalues of  $f(A)$  are  $f(\lambda_1), \dots, f(\lambda_n)$ . Since  $f(A)$  is a positive matrix, the Perron-Frobenius theorem [7, p.123] ensures that the spectral radius of  $f(A)$  is a simple eigenvalue (Perron root). Suppose  $f(\lambda_j)$  is the Perron root of  $f(A)$ . Then  $\lambda_j$  is a simple eigenvalue of  $A$ , for otherwise  $f(\lambda_j)$  would be a multiple eigenvalue of  $f(A)$ . We claim that  $\lambda_j$  is a real number. To see the claim, suppose to the contrary that  $\lambda_j$  is nonreal. Since the nonreal eigenvalues of any real matrix occur in conjugate pairs,  $\overline{\lambda_j}$  is also an eigenvalue of  $A$  and  $\overline{\lambda_j} \neq \lambda_j$ , where the bar notation means the complex conjugate. Since  $f(\lambda_j)$  is a real number and  $f$  is a real polynomial,  $f(\lambda_j) = \overline{f(\lambda_j)} = f(\overline{\lambda_j})$  which shows that  $f(\lambda_j)$  is a multiple eigenvalue of  $f(A)$ , a contradiction. Thus  $\lambda_j$  is a simple real eigenvalue of  $A$ .

Every left or right eigenvector of  $A$  corresponding to  $\lambda_j$  is also a left or right eigenvector of  $f(A)$  corresponding to the Perron root  $f(\lambda_j)$ . Since  $f(\lambda_j)$  is simple, it has geometric multiplicity 1. Hence every left or right eigenvector of  $A$  corresponding to  $\lambda_j$  is a real scalar multiple of a left or right Perron vector of  $f(A)$ . It follows that  $A$  has positive left and right eigenvectors corresponding to the simple real eigenvalue  $\lambda_j$ .

Conversely, let  $A$  be a real matrix of order  $n$  and suppose that  $A$  has a simple real eigenvalue  $r$  with corresponding left and right positive eigenvectors  $v^T$  and  $u$  respectively. Without loss of generality we normalize  $u, v$  so that  $v^T u = 1$ . Let  $q$  be the minimal polynomial of  $A$ , and let  $p(x) = q(x)/(x - r)$ . Then  $0 = q(A) = p(A)(A - rI)$ . Applying Sylvester's inequality on the rank of a matrix product and using the condition that  $\text{rank}(A - rI) = n - 1$ , we have  $\text{rank } p(A) \leq 1$ . Since  $q$  is the minimal polynomial and

$\deg p = (\deg q) - 1$ , we have  $p(A) \neq 0$ . Hence  $\text{rank } p(A) = 1$ . Since  $r$  is a simple eigenvalue,  $p(r) \neq 0$ . Now it is easy to show that the conditions

$$\text{rank } p(A) = 1, \quad p(A)u = p(r)u, \quad v^T p(A) = p(r)v^T, \quad p(r) \neq 0, \quad v^T u = 1$$

imply that  $p(A) = p(r)uv^T$ . Since  $u$  and  $v$  are positive vectors, we have

$$(p(A))^2 = (p(r))^2 uv^T > 0.$$

This proves that  $A$  is algebraically positive.  $\square$

From the above proof of Theorem 1 we see that a real matrix  $A$  is algebraically positive if and only if there is a real polynomial  $f$  such that  $f(A)$  is irreducible and nonnegative. This fact also follows from the result [7, p.121] that if  $B$  is an irreducible nonnegative matrix of order  $n$ , then  $(I + B)^{n-1} > 0$ .

**Remark 2** *Algebraically positive matrices have the following properties.*

(i) *Every algebraically positive matrix is irreducible.*

(ii) *If  $A$  is a real matrix and there is a positive integer  $k$  such that  $A^k$  is algebraically positive, then so is  $A$ .*

**Proof.** (i) If  $A$  is a reducible real matrix, then there is a permutation matrix  $P$  such that  $A = P^T \begin{bmatrix} A_1 & 0 \\ A_2 & A_3 \end{bmatrix} P$  where  $A_1$  and  $A_3$  are square and nonvoid. For any real polynomial  $f$  we have  $f(A) = P^T \begin{bmatrix} f(A_1) & 0 \\ \star & f(A_3) \end{bmatrix} P$ . Thus  $f(A)$  has zero entries and cannot be positive.

(ii) Obvious by definition.  $\square$

It is straightforward to see from the definition that if a matrix  $A$  is algebraically positive, then for each  $t \in \mathbb{R}$ ,  $A + tI$  is also algebraically positive. Hence, shifting the entire diagonal of an algebraically positive matrix will preserve that property. However, as the following example shows, perturbing a single diagonal entry may not preserve algebraic positivity.

**Example 3** Consider the matrices

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 2 & -1 & 1 \\ 2 & -2 & -1 \end{bmatrix}, \quad B = \begin{bmatrix} -2 & 2 & 1 \\ 2 & -1 & 1 \\ 2 & -2 & -1 \end{bmatrix}, \quad (1)$$

which differ only in the first diagonal entry. Since

$$2I + 2A + A^2 = \begin{bmatrix} 16 & 4 & 5 \\ 8 & 3 & 2 \\ 2 & 4 & 1 \end{bmatrix} > 0,$$

we see that  $A$  is algebraically positive. Let  $f(x) = a + bx + cx^2$  be a real polynomial. Consider the entries in the positions  $(1, 2)$  and  $(3, 2)$  of

$$f(B) = aI + b \begin{bmatrix} -2 & 2 & 1 \\ 2 & -1 & 1 \\ 2 & -2 & -1 \end{bmatrix} + c \begin{bmatrix} 10 & -8 & -1 \\ -4 & 3 & 0 \\ -10 & 8 & 1 \end{bmatrix}.$$

To make the  $(1, 2)$  entry positive we must have  $b > 4c$ . But to make the  $(3, 2)$  entry positive we must have  $b < 4c$ . Thus no  $a, b, c$  can make  $f(B)$  positive. This shows that  $B$  is not algebraically positive.

We remark that irreducibility and having a simple real eigenvalue are not sufficient for algebraic positivity. The matrix  $B$  in Example 3 is irreducible and has three simple real eigenvalues  $0, -2 + \sqrt{3}, -2 - \sqrt{3}$ .

Recall that a matrix  $A$  is said to be *skew-symmetric* if  $A^T = -A$ .

**Corollary 4** *No nonsingular skew-symmetric real matrix is algebraically positive.*

**Proof.** Since all the eigenvalues of any skew-symmetric real matrix are pure imaginary, the only possible real eigenvalue is 0. Hence a nonsingular skew-symmetric real matrix has no real eigenvalue and is not algebraically positive by Theorem 1.  $\square$

Note that the nonsingularity condition in Corollary 4 cannot be dropped. For example,

$$A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}, \quad 3I + A^2 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} > 0$$

showing that the skew-symmetric real matrix  $A$  is algebraically positive. Of course  $A$  is singular.

**Theorem 5** *If  $A$  is an irreducible real matrix all of whose off-diagonal entries are non-negative (or nonpositive), then  $A$  is algebraically positive.*

**Proof.** It suffices to consider the nonnegative case by considering  $-A$  if necessary. Choose a positive number  $d$  such that  $d$  is larger than the maximum absolute value of the diagonal entries of  $A$ . Then  $dI + A$  is an irreducible nonnegative matrix. It follows [7, p.121] that  $(I + (dI + A))^{n-1} > 0$  where  $n$  is the order of  $A$ ; i.e.,  $((d+1)I + A)^{n-1} > 0$ . Thus  $A$  is algebraically positive.  $\square$

**Corollary 6** *A nonnegative matrix is algebraically positive if and only if it is irreducible.*

**Proof.** This follows immediately from Remark 2(i) and Theorem 5.  $\square$

The next example shows that similarity transformations may not preserve algebraic positivity.

**Example 7** Consider the matrix  $A$  given by

$$A = - \begin{bmatrix} 2 & 1 & 1 \\ 2 & 1 & 1 \\ 0 & 2 & 1 \end{bmatrix}$$

and note that  $A$  is algebraically positive by Theorem 5. Let

$$B = T^{-1}AT = \begin{bmatrix} -2 & 2 & 1 \\ 2 & -1 & 1 \\ 2 & -2 & -1 \end{bmatrix} \quad \text{where } T = \begin{bmatrix} 1 & -1 & 0 \\ 0 & -1 & -1 \\ 2 & 0 & 1 \end{bmatrix}.$$

Then  $B$  is similar to  $A$ . Note that this matrix  $B$  is the same  $B$  as in Example 3 where we have proved that it is not algebraically positive.

**Definition.** If a real matrix  $A$  is algebraically positive, we define the *index of algebraic positivity* of  $A$  to be the least degree of a real polynomial  $f$  such that  $f(A)$  is positive.

Clearly, this index of an algebraically positive matrix of order  $n \geq 2$  is an integer between 1 and  $n - 1$ . Every real matrix of order 1 is algebraically positive and has index of algebraic positivity 0.

Recall that for a primitive nonnegative matrix  $A$ , the *exponent* of  $A$  is the smallest positive integer  $m$  such that  $A^m > 0$  (see [2, p.78]). For a nonnegative matrix of order  $n$ , the exponent must lie in the interval  $[1, n^2 - 2n + 2]$ , and it is well-known ([2, p.84-85], [8]) that the exponents of primitive matrices of a given order ( $n \geq 4$ ) have gaps – i.e. there

are integers  $k \in [1, n^2 - 2n + 2]$  that cannot be realized as the exponent of any primitive  $n \times n$  matrix. The following result shows that the indices of algebraically positive matrices of a given order have no gaps.

**Theorem 8** *Let  $n, k$  be integers such that  $1 \leq k \leq n-1$ . Then there exists an algebraically positive matrix of order  $n$  whose index of algebraic positivity is equal to  $k$ .*

**Proof.** The assumption  $1 \leq k \leq n-1$  implies that  $n \geq 2$ . Any positive matrix of order at least 2 has index of algebraic positivity 1. Next suppose  $k \geq 2$ . Let  $A$  be the adjacency matrix of the graph  $T$  which is a tree with vertices  $1, 2, \dots, n$  consisting of the  $(1, k)$ -path with vertices  $1, 2, \dots, k$  and the edges  $ki, i = k+1, k+2, \dots, n$ . Then  $A$  is of order  $n$ . Note that there is a loop at every vertex of the digraph of the matrix  $I + A$ . Since the diameter of  $T$  is  $k$ , it follows that  $(I + A)^k > 0$  and for any real polynomial  $f$  of degree  $\leq k-1$ ,  $f(A)$  has zero entries. In fact, the entry of  $f(A)$  in the position  $(1, n)$  is zero, since in the digraph of  $A$  there is no walk from vertex 1 to vertex  $n$  of length  $\leq k-1$ . Hence the index of algebraic positivity of  $A$  is equal to  $k$ .  $\square$

If  $A$  is an irreducible nonnegative matrix with index of imprimitivity  $k \geq 2$ , then  $A$  is permutationally similar to a matrix of the form

$$\begin{bmatrix} 0 & A_{12} & 0 & \cdots & 0 \\ 0 & 0 & A_{23} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A_{k-1,k} \\ A_{k1} & 0 & 0 & \cdots & 0 \end{bmatrix} \quad (2)$$

where every zero block on the diagonal is a square matrix [7, p.136]. The matrix in (2) is called the *cyclic canonical form* of  $A$ .

**Theorem 9** *Let  $A$  be an irreducible nonnegative matrix of order  $n$  and let  $\alpha$  and  $k$  be the index of algebraic positivity and the index of imprimitivity of  $A$  respectively. If  $k \leq n-1$  then*

$$\alpha \geq k. \quad (3)$$

When  $k = 1$ , equality in (3) holds if and only if all the off-diagonal entries of  $A$  are positive. When  $2 \leq k \leq n-1$  and the cyclic canonical form of  $A$  is the matrix in (2), equality in (3) holds if and only if all the matrices  $A_{12}, A_{23}, \dots, A_{k-1,k}, A_{k1}$  are positive. If  $k = n$ , then  $\alpha = n-1$ .

**Proof.** First note that  $A$  is algebraically positive by Corollary 6. The case when  $k = 1$  is clear. Now suppose  $2 \leq k \leq n - 1$ . By a permutation similarity transformation if necessary, it suffices to consider the case when  $A$  itself is the matrix in (2) and we make this assumption. The condition  $k \leq n - 1$  implies that at least one zero block on the diagonal of  $A$  has order  $\geq 2$ . Since the  $p$ -th powers of  $A$  for  $p = 1, 2, \dots, k$  are

$$A = \begin{bmatrix} 0 & A_{12} & 0 & \cdots & 0 \\ 0 & 0 & A_{23} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A_{k-1,k} \\ A_{k1} & 0 & 0 & \cdots & 0 \end{bmatrix}, A^2 = \begin{bmatrix} 0 & 0 & A_{12}A_{23} & 0 & \cdots & 0 \\ 0 & 0 & 0 & A_{23}A_{34} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ A_{k-1,k}A_{k1} & 0 & 0 & 0 & \cdots & 0 \\ 0 & A_{k1}A_{12} & 0 & 0 & \cdots & 0 \end{bmatrix},$$

$$\dots, A^k = \begin{bmatrix} A_{12}A_{23} \cdots A_{k1} & 0 & 0 & \cdots & 0 \\ 0 & A_{23} \cdots A_{k1}A_{12} & 0 & \cdots & 0 \\ 0 & 0 & A_{34} \cdots A_{12}A_{23} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A_{k1}A_{12} \cdots A_{k-1,k} \end{bmatrix},$$

for any real polynomial  $f$  of degree  $\leq k - 1$ ,  $f(A)$  has zero entries in some diagonal block and hence is not positive. This shows the inequality (3).

If all the matrices  $A_{12}, A_{23}, \dots, A_{k-1,k}, A_{k1}$  are positive, we have  $A + A^2 + \dots + A^k > 0$ . Hence  $\alpha \leq k$  which, together with  $\alpha \geq k$ , yields  $\alpha = k$ . Conversely suppose  $\alpha = k$ . Then there are real numbers  $c_0, c_1, \dots, c_k$  such that  $c_0I + c_1A + \dots + c_kA^k > 0$ . Since  $A_{12}, A_{23}, \dots, A_{k-1,k}, A_{k1}$  and their cyclic products appearing in  $A, A^2, \dots, A^k$  lie in different positions, we must have that  $A_{12}, A_{23}, \dots, A_{k-1,k}, A_{k1}$  are all positive.

If  $k = n$ , each of the matrices  $A_{12}, A_{23}, \dots, A_{k-1,k}, A_{k1}$  must be of order 1, and hence is a positive number, since  $A$  is irreducible. For any real polynomial  $f$  of degree  $\leq n - 2$ ,  $f(A)$  has zero entries, showing that  $\alpha \geq n - 1$ . On the other hand, since the order of  $A$  is  $n$ , we always have  $\alpha \leq n - 1$ . Thus  $\alpha = n - 1$ .  $\square$

**Remark** The idea in the proof of Theorem 9 can be used to show that if the digraph of a real matrix  $A$  is a Hamilton cycle and  $A$  has both positive and negative entries, then  $A$  is not algebraically positive.



### 3 Algebraic positivity and sign patterns

A *sign pattern* is a matrix whose entries are from the set  $\{+, -, 0\}$ . The *sign pattern of a real matrix*  $A$  is the matrix obtained from  $A$  by replacing each entry by its sign. Whether a nonnegative matrix is primitive depends only on its zero–nonzero pattern [2, Section 3.4], so that primitivity is a purely combinatorial property. This is not the case for algebraic positivity: the following example shows that the sign pattern alone may not determine whether or not a real matrix is algebraically positive.

**Example 10** Let

$$A = \begin{bmatrix} 0 & 30 & -1 & -5 \\ -1 & 0 & 40 & -5 \\ 50 & -3 & 0 & 60 \\ 10 & 60 & -1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & -1 & -1 \\ -1 & 0 & 1 & -1 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & 0 \end{bmatrix}, \quad C = \begin{bmatrix} 0 & + & - & - \\ - & 0 & + & - \\ + & - & 0 & + \\ + & + & - & 0 \end{bmatrix}.$$

The two matrices  $A$  and  $B$  have the same sign pattern  $C$ .

$$1000A + 20A^2 + A^3 = \begin{bmatrix} 55847 & 3945 & 11560 & 65235 \\ 62800 & 193782 & 17765 & 35960 \\ 43110 & 103680 & 199032 & 12985 \\ 122897 & 30990 & 59440 & 134235 \end{bmatrix} > 0$$

shows that  $A$  is algebraically positive. Since  $B$  is a nonsingular skew-symmetric real matrix ( $\det B = 1$ ),  $B$  is not algebraically positive by Corollary 4.

Given a sign pattern  $A$ , the *pattern class* of  $A$ , denoted  $Q(A)$ , is the set of real matrices whose sign patterns are  $A$ . Let  $P$  be a property about real matrices. A sign pattern  $A$  is said to *require*  $P$  if every matrix in  $Q(A)$  has property  $P$ ;  $A$  is said to *allow*  $P$  if there exists a matrix in  $Q(A)$  that has property  $P$ . For example, let

$$A = \begin{bmatrix} 0 & + & - \\ - & 0 & + \\ + & - & 0 \end{bmatrix}, \quad B = \begin{bmatrix} + & + & + \\ + & - & + \\ + & - & - \end{bmatrix}, \quad C = \begin{bmatrix} 0 & + & 0 \\ - & 0 & + \\ 0 & - & 0 \end{bmatrix}.$$

Then  $A$  requires algebraic positivity (since for any  $M \in Q(A)$ ,  $M^2$  has all positive off-diagonal entries),  $B$  allows algebraic positivity (since  $\begin{bmatrix} 3 & 3 & 3 \\ 3 & -1 & 3 \\ 3 & -1 & -1 \end{bmatrix} \in Q(B)$  and has

a positive square), and  $C$  does not allow algebraic positivity (since no matrix in  $Q(C)$  admits a positive right eigenvector).

A sign pattern  $A$  of order  $n$  is said to be *spectrally arbitrary* if given any monic real polynomial  $f$  of degree  $n$ , there is a matrix in  $Q(A)$  whose characteristic polynomial is  $f$  [3]. We have the following (negative) result.

**Theorem 11** *No spectrally arbitrary sign pattern requires algebraic positivity.*

**Proof.** It is clear that any sign pattern of order 1 is not spectrally arbitrary. Let  $A$  be a spectrally arbitrary sign pattern of order  $n \geq 2$ . Then there is a real matrix  $B$  in  $Q(A)$  whose characteristic polynomial is  $f(x) = (x - 1)^n$ . Obviously  $B$  has no simple eigenvalue. By Theorem 1,  $B$  is not algebraically positive. This shows that  $A$  does not require algebraic positivity.  $\square$

Next we give a necessary condition for a sign pattern to allow algebraic positivity.

**Theorem 12** *If a sign pattern allows algebraic positivity, then every row and column contains a +, or every row and column contains a -.*

**Proof.** Let  $A$  be a sign pattern that allows algebraic positivity and suppose that  $M \in Q(A)$  is an algebraically positive matrix. By Theorem 1, there is a real eigenvalue  $r$ , a positive right eigenvector  $v$ , and a positive left eigenvector  $w^T$  such that  $Mv = rv$  and  $w^T M = rw^T$ . If  $r > 0$ , then each row of  $M$  contains a positive entry (otherwise we have a contradiction to the fact that  $Mv = rv$ ) and similarly every column of  $M$  contains a positive entry. If  $r < 0$ , an analogous argument shows that every row and column of  $M$  must contain a negative entry. Finally, if  $r = 0$ , then since  $Mv = 0$ , each row must contain both a negative entry and a positive entry (recall that  $M$  is irreducible and so contains no all zero row or column) and similarly since  $w^T M = 0^T$ , each column of  $M$  contains both a positive entry and a negative entry.  $\square$

The following example shows that the necessary condition of Theorem 12 is not sufficient for a sign pattern to allow algebraic positivity.

**Example 13** Consider the sign pattern  $A = \begin{bmatrix} + & + \\ - & + \end{bmatrix}$ . Observe that if  $M \in Q(A)$ , then

$M$  has the form  $\begin{bmatrix} a & b \\ -c & d \end{bmatrix}$  for some  $a, b, c, d > 0$ . Observe that for any  $x, y \in \mathbb{R}$ ,  $xI + yM$  has a nonpositive off-diagonal entry. Hence,  $M$  cannot be algebraically positive.

In the special case of symmetric tridiagonal sign patterns, we have the following characterization of the allow problem for algebraic positivity.

**Theorem 14** *Suppose that  $A$  is an irreducible  $n \times n$  symmetric tridiagonal sign pattern matrix. Then  $A$  allows algebraic positivity if and only if either every row of  $A$  contains a  $+$ , or every row of  $A$  contains a  $-$ .*

**Proof.** If  $A$  allows algebraic positivity, then the condition on the rows follows immediately from Theorem 12. Suppose now for concreteness that every row of  $A$  contains a  $+$ . For each  $i = 1, \dots, n$ , denote the number of  $+$  and  $-$  entries in row  $i$  of  $A$  by  $n_+(i)$  and  $n_-(i)$ , respectively. Construct a matrix  $M \in Q(A)$  as follows: put a  $-1$  in the positions of  $M$  corresponding to  $-$  entries in  $A$ , and for each  $+$  entry in row  $i$  of  $A$ , put a  $\frac{1+n_-(i)}{n_+(i)}$  in the corresponding position of  $M$ . Since the row sums of  $M$  are all 1, the all ones vector is a right eigenvector of  $M$  corresponding to the eigenvalue 1. Since  $M$  is irreducible, sign-symmetric and tridiagonal, every eigenvalue of  $M$  is real and simple. Further, there is a diagonal matrix  $D$  with positive entries on the diagonal such that  $\hat{M} = DMD^{-1}$  is symmetric. Evidently  $\text{diag}(D)$  is a positive right eigenvector of  $\hat{M}$  corresponding to the eigenvalue 1. Hence  $\hat{M}$  has 1 as a simple eigenvalue with corresponding left and right eigenvectors that are positive. Consequently  $\hat{M}$  is algebraically positive by Theorem 1.  $\square$

We now characterize a special class of  $3 \times 3$  sign patterns that require algebraic positivity.

**Theorem 15** *Suppose that  $A$  is an irreducible  $3 \times 3$  symmetric tridiagonal sign pattern matrix. Then  $A$  requires algebraic positivity if and only if one of the following holds:*

- i) all nonzero off-diagonal entries of  $A$  are  $+$ ;*
- ii) all nonzero off-diagonal entries of  $A$  are  $-$ ;*
- iii)  $A$  is permutationally similar to a matrix in the set*

$$S = \left\{ \begin{bmatrix} 0 & + & 0 \\ + & 0 & - \\ 0 & - & + \end{bmatrix}, \begin{bmatrix} - & + & 0 \\ + & 0 & - \\ 0 & - & + \end{bmatrix}, \begin{bmatrix} 0 & + & 0 \\ + & + & - \\ 0 & - & + \end{bmatrix}, \right. \\ \left. \begin{bmatrix} - & + & 0 \\ + & + & - \\ 0 & - & + \end{bmatrix}, \begin{bmatrix} 0 & + & 0 \\ + & - & - \\ 0 & - & + \end{bmatrix}, \begin{bmatrix} - & + & 0 \\ + & - & - \\ 0 & - & + \end{bmatrix} \right\};$$

- iv)  $-A$  is permutationally similar to a matrix in  $S$ .*

**Proof.** Assume that  $A$  requires algebraic positivity. If all off-diagonal entries of  $A$  have the same sign, then either i) or ii) holds. Suppose now that  $A$  has two off-diagonal entries  $+$  and two off-diagonal entries  $-$ . By considering  $-A$  if necessary, we may assume by Theorem 12 that every row of  $A$  contains a  $+$ . Further, simultaneously permuting rows and columns of  $A$  if necessary, we may further assume that the  $(1, 2)$  and  $(2, 1)$  entries of  $A$  are  $+$ , and the  $(2, 3)$  and  $(3, 2)$  entries are  $-$ . Since the  $(3, 1)$  entry of  $A$  is 0, we find from Theorem 12 that necessarily the  $(3, 3)$  entry of  $A$  is  $+$ .

Let  $M \in Q(A)$ ; without loss of generality we may take  $M$  to be symmetric, and multiplying  $M$  by a suitable positive scalar if necessary, we take  $M$  to have the form  $M = \begin{bmatrix} a & 1 & 0 \\ 1 & b & -c \\ 0 & -c & d \end{bmatrix}$ , where  $c, d > 0$ . Observe that  $M^2 = \begin{bmatrix} a^2 + 1 & a + b & -c \\ a + b & 1 + b^2 + c^2 & -c(b + d) \\ -c & -c(b + d) & c^2 + d^2 \end{bmatrix}$ . Let  $p(x) = \alpha x^2 + \beta x + \gamma$  be a polynomial such that  $p(M) > 0$ . By considering the  $(1, 3)$  entry of  $p(M)$ , we deduce that  $\alpha < 0$ . Considering the  $(1, 2)$  and  $(2, 3)$  entries of  $p(M)$  now yields that  $\beta - |\alpha|(a + b) > 0$  and  $-\beta c + |\alpha|(b + d)c > 0$ . Setting  $\sigma = \frac{\beta}{|\alpha|}$  and simplifying, the preceding inequalities become  $\sigma > a + b$  and  $b + d > \sigma$ . In particular it must be the case that  $d > a$ . Suppose that the  $(1, 1)$  entry of  $A$  is  $+$ . Then we may find an  $M \in Q(A)$  such that  $m_{1,1} = 1$  and  $m_{3,3} = 1$ , contradicting the fact that  $d > a$  above. We thus conclude that the  $(1, 1)$  entry of  $A$  must be either a 0 or a  $-$ . Observe that the sign patterns in the set  $S$  correspond to the six cases arising from the  $(1, 1)$  entry of  $A$  being 0 or  $-$ , and the  $(2, 2)$  entry being 0,  $+$  or  $-$ . Hence  $A \in S$ .

Next suppose that  $A$  satisfies one of i)–iv). If either i) or ii) is satisfied, then  $A$  requires algebraic positivity by Theorem 5. Suppose now that  $A$  satisfies iii). As above, if  $M \in Q(A)$ , then without loss of generality, we may take  $M$  to have the form  $M = \begin{bmatrix} a & 1 & 0 \\ 1 & b & -c \\ 0 & -c & d \end{bmatrix}$ , where  $c, d > 0$  and  $a \leq 0$ . Set  $\sigma = b + \frac{a+d}{2}$ , and note that

$$-M^2 + \sigma M = \begin{bmatrix} \sigma a - a^2 - 1 & \sigma - a - b & c \\ \sigma - a - b & \sigma b - 1 - b^2 - c^2 & c(b + d) - \sigma c \\ c & c(b + d) - \sigma c & \sigma d - c^2 - d^2 \end{bmatrix} = \begin{bmatrix} \sigma a - a^2 - 1 & \frac{d-a}{2} & c \\ \frac{d-a}{2} & \sigma b - 1 - b^2 - c^2 & c(\frac{d-a}{2}) \\ c & c(\frac{d-a}{2}) & \sigma d - c^2 - d^2 \end{bmatrix},$$

which has positive off-diagonal entries since  $d > 0 \geq a$  and  $c > 0$ . Adding a suitable multiple of the identity matrix now shows that  $-M^2 + \sigma M + tI > 0$  for some  $t \in \mathbb{R}$ , so that  $A$  requires algebraic positivity.

Finally, if  $A$  satisfies iv), it immediately follows from iii) that  $A$  requires algebraic positivity.  $\square$

We close the paper with the following problems.

**Problem 1.** Characterize those sign patterns that require algebraic positivity.

**Problem 2.** Characterize those sign patterns that allow algebraic positivity.

The approach and results in [5] might be helpful in studying these problems.

*Acknowledgement.* The authors are grateful to two anonymous referees whose comments helped to improve the presentation of the results in this paper.

## References

- [1] R. Bhatia, Matrix Analysis, GTM 169, Springer, New York, 1997.
- [2] R.A. Brualdi and H.J. Ryser, Combinatorial Matrix Theory, Cambridge University Press, New York, 1991.
- [3] J.H. Drew, C.R. Johnson, D.D. Olesky and P. van den Driessche, Spectrally arbitrary patterns, Linear Algebra Appl., 308(2000), 121-137.
- [4] R.A. Horn and C.R. Johnson, Topics in Matrix Analysis, Cambridge University Press, Cambridge, 1991.
- [5] S.J. Kirkland, J.J. McDonald and M.J. Tsatsomeros, Sign-patterns which require a positive eigenvalue, Linear and Multilinear Algebra, 41(1996), 199-210.
- [6] J.J. McDonald, P. Paparella and M.J. Tsatsomeros, Matrix roots of eventually positive matrices, Linear Algebra Appl., 456(2014), 122-137.
- [7] X. Zhan, Matrix Theory, GSM 147, Amer. Math. Soc., Providence, RI, 2013.
- [8] K.M. Zhang, On Lewin and Vitek's conjecture about the exponent set of primitive matrices, Linear Algebra Appl., 96(1987), 101-108.