

Hyperbolic Groups (Presentation Notes)

Dana Kapoostinsky

March 2024

1 Motivation

Why are hyperbolic groups interesting?

- From a probabilistic point of view, most¹ finitely generated groups are hyperbolic.
- Hyperbolic groups have solvable word problem².
- As a result, “most” finitely generated groups have solvable word problem.

We use geometric techniques to prove the second point.

¹The underlying statistical model requires that the generating set $\{a_1, \dots, a_n\}$ and the number of relators be fixed. Next, set an upper bound l on the word length of each relator. Finally, for each relator, choose a word at random (uniformly and independently) from the set of reduced words over $\{a_1, \dots, a_n\}$ of length at most l . [1]

²Given a group $G = \langle S | R \rangle$ and an arbitrary word $w \in S^*$, is $w =_G 1$? The word problem is undecidable.

2 Cayley Graphs

Intuitively: A Cayley graph is a graph that models the multiplication of a group G .

- Vertices are labelled by elements of G .
- Edges are labelled by elements of the generating set S of G .

Example. Here are two distinct Cayley graphs for the group $(\mathbb{Z}, +)$.




Figure 1: The Cayley graph $\text{Cay}_{\{1\}}(\mathbb{Z})$.



Figure 2: The Cayley graph $\text{Cay}_{\{1,2\}}(\mathbb{Z})$. (Notice how the redundant generator added “extra” edges to the graph.)

Notice that $\text{Cay}_{\{1\}}(\mathbb{Z})$ and $\text{Cay}_{\{1,2\}}(\mathbb{Z})$ are *not* isomorphic as graphs. So, Cayley graphs are not unique.

3 Geodesic Metric Spaces

We want to associate a group G to a metric space. We do this by metrizing $X = \text{Cay}_S(G)$. In particular, it is desirable to turn X into a geodesic metric space.

Definition. Let (X, d) be a metric space and let $L \in \mathbb{R}_{\geq 0}$. A path p between points x, y in X is geodesic if

1. there exists an isometric embedding from $[0, L] \rightarrow p$ and
2. if p is the shortest path between x and y .

Definition. A space (X, d) is geodesic if for any pair of points $x, y \in X$, there exists a geodesic path between x and y .

Metrizing $X = \text{Cay}_S(G)$:

- Identify each edge in X with the unit interval $[0, 1]$.
- Define the distance between two vertices x and y to be the length of the shortest edge path connecting x and y .³


We now have a way to associate a geodesic metric space to any (finitely generated) group G .

³This is commonly referred to as the path or word metric.

4 Hyperbolic Groups

Definition. A triangle xyz is geodesic if $\overline{xy}, \overline{yz}, \overline{xz}$ are geodesic paths.

Definition. Let $\delta \in \mathbb{R}_{\geq 0}$. A triangle xyz is δ -slim if the δ nbhd of \overline{xy} and the δ nbhd of \overline{xz} cover all of xyz .



Definition. A geodesic metric space (X, d) is hyperbolic (or δ -hyperbolic) if there exists $\delta \in \mathbb{R}_{\geq 0}$ such that all geodesic triangles in X are δ -slim.




Figure 3: Geodesic triangles form tripods in trees.




Figure 4: Euclidean space is *not* δ -hyperbolic.

Definition. A group G is hyperbolic if its cayley graph $\text{Cay}_S(G)$ is δ -hyperbolic.

(Note: Implicit in this definition is that G is necessarily finitely generated if it is hyperbolic.)

A PROBLEM WITH THIS DEFINITION:

- We already know that Cayley graphs are *not* unique!
- Given two distinct cayley graphs X and X' of G , what if X is hyperbolic but X' is not?

It turns out that this second point can't happen, so we have a well-defined definition. Let's see why...

5 Quasi-Isometric Embeddings

Definition. Let (M_1, d_1) and (M_2, d_2) be two metric spaces and let $f : M_1 \rightarrow M_2$. Then f is a quasi-isometric embedding if there exist $K, C \in \mathbb{R}^+$ such that

$$\frac{1}{K}d_2(f(x), f(y)) - C \leq d_1(x, y) \leq Kd_2(f(x), f(y)) + C$$

for all $x, y \in M_1$.

Quasi-isometric embeddings turn out to be a very important tool in geometric group theory. Essentially, we want to study properties that are invariant under QI embeddings.

Intuitively:

- Quasi-isometric embeddings are like isometric embeddings with some tolerance for “error” on a small enough scale, *i.e.*, quasi-isometries respect large scale geometry.
- If $X \sim_{QI} Y$, then if we zoom out far enough these two spaces should “look” the same.

Examples:

- Recall the two Cayley graphs given earlier in figures 1 and 2.
- Set $M_1 = \text{Cay}_{\{1\}}(\mathbb{Z})$ and $M_2 = \text{Cay}_{\{1,2\}}(\mathbb{Z})$.
- Then taking $f : M_1 \rightarrow M_2$ to be an embedding with $K = 1$ and $C = 0$ is a quasi-isometric embedding.
- Or, define $g : M_2 \rightarrow M_1$ by mapping black edges to black edges and a blue edge to its neighbouring black edges in the obvious way. Take $K = 2$ and $C = 0$. This is also a QI embedding.

Some other examples:

- All finite graphs (with path metric) are quasi-isometric.
- \mathbb{R} and \mathbb{Z} are quasi-isometric via the map $f : \mathbb{R} \rightarrow \mathbb{Z}$ defined by $f : x \mapsto \lfloor x \rfloor$.
- \mathbb{Z} and \mathbb{Z}^2 are *not* quasi-isometric (idea of proof: asymptotic argument... show that balls in \mathbb{Z}^2 grow much faster than those in \mathbb{Z} , so there cannot exist values K, C that would allow you to set a bound on the corresponding distances).

Theorem. [2] Let X and X' be two distinct Cayley graphs for a group G . Then $X \sim_{QI} X'$.

Theorem. [2] If $X \sim_{QI} X'$, then X is hyperbolic if and only if X' is hyperbolic.

The proofs for both of these theorems require very careful consideration of δ -slim triangles and QI embeddings. (Very LONG and cumbersome chains of inequalities!)

Now we have a sensible definition for what it means for a group to be hyperbolic. The next goal is to show that hyperbolic groups have Dehn presentation using a geometric argument.

6 Dehn Presentation

Definition. Let S be a finite alphabet and $n \in \mathbb{N}$. The group presentation

$$\langle S \mid u_1 v_1^{-1} = \cdots = u_n v_n^{-1} = 1 \rangle$$

is a Dehn presentation if:

- For all $1 \leq i \leq n$, the length of the word v_i is shorter than the length of the word u_i , and
- Any freely reduced word non empty w over $S \cup S^{-1}$ such that $w =_G 1$ must contain a subword of the form u_i or u_i^{-1} .

Groups with Dehn presentation have solvable word problem.

Shortening algorithm:

- Choose a word $w \in G$.
- Freely reduce w .
- Check to see if w contains a subword of the form u_i or u_i^{-1} . (This can always be done because w is finite.) If no, then $w \neq_G 1$.
- If yes, replace the occurrence of u_i with v_i . Then $w = w' v_i w''$. Repeat procedure on the word $w' v_i w''$. This always terminates because replacing u_i with v_i reduces the length of the word.

Finally, given a hyperbolic group G we construct a Dehn presentation for G by following non-geodesic paths in $\text{Cay}_S(G)$.

7 Taming Quasi-Geodesics

Think of a “non-geodesic” path as the image of some quasi-isometric embedding $\gamma : [0, L] \rightarrow X$. Such a path is commonly referred to as a quasi-geodesic.

We are interested in a special kind of quasi-geodesic path:

Definition. A path p in X is k -local geodesic if every subpath q of p with $|q| \leq k$ is geodesic.


Intuitively: k -local geodesic paths are almost geodesic (they are geodesic on a small enough scale).

The following proof offers one example of how the geometry of δ -slim triangles can be used to obtain a “nice” property of hyperbolic spaces.

Lemma 1. [3] Let X be δ -hyperbolic. Set $k > 8\delta$. Suppose p is a k -local geodesic path in X . Then there exists a geodesic path q sharing the same end points as p such that p is contained in a 2δ nbhd of q .

In other words, in hyperbolic spaces, k -local geodesics stay close to geodesics.

Proof. First, note that the path p can be described by the image of a quasi-isometric embedding $\gamma : [0, L] \rightarrow X$.



Choose M to be a point on p having maximal distance from q . Then there exists some $t_M \in [0, L]$ such that $\gamma(t_M) = M$.


Either:

1. $d(0, t_M) < 4\delta$ or
2. $d(0, t_M) > 4\delta$ (and $d(t_M, L) > 4\delta$).

Case 1: Suppose $d(0, t_M) < 4\delta$. Fix a point A on p such that

$$d_X(M, A) > 4\delta,$$


and $p|_{I,A}$ is geodesic.⁴



Note: This is possible because p must have length greater than 8δ since it's a k -local geodesic.

Now, choose a point B on q with minimal distance from A .


⁴Notation: $p|_{I,A}$ denotes the subarc of p with end points $I = \gamma(0)$ and A .



Consider the triangle $\triangle IAB$. This is a geodesic triangle. So, there exists a point x on either \overline{IB} or \overline{AB} such that

$$d_X(M, x) \leq \delta$$

(by the definition of δ -slim triangle).



Suppose $x \in \overline{AB}$, then

$$\begin{aligned} d_X(M, x) - d_X(A, x) &= d_X(M, A) \\ &> 4\delta && \text{by previous assumption} \\ \implies d_X(A, x) &> 3\delta && (*) \end{aligned}$$

Also,

$$\begin{aligned}d_X(M, B) - d_X(A, B) &\leq (d_X(M, x) + d_X(x, B)) - (d_X(A, x) + d_X(x, B)) \\ &= d_X(M, x) - d_X(A, x) \\ &< \delta - 3\delta && \text{by } (\star) \\ &< 0,\end{aligned}$$

which contradicts the fact that M was chosen to have maximal distance from the path q . So, x must lie on \overline{IA} .

Now, for any other point $y \in p$,

$$d_X(y, x) \leq d_X(M, x) \leq \delta.$$

Therefore, p is contained in a δ -nbhd of q .

The proof for case 2 is omitted. For complete proof, see [3] □

Lemma 2. [2] Let X be a δ -hyperbolic space. Any closed loop γ in X contains a subarc p such that $|p| \leq 8\delta$ and p is *not* geodesic.

Proof. By contradiction. Use lemma 1.

Theorem. [4] If G is hyperbolic, then G admits a Dehn presentation.

Proof. We can assume G has finite generating set S and $X = \text{Cay}_S(G)$ is δ -hyperbolic.

Step 1: Define a procedure to construct a set of Dehn relators:

- Fix $k > 8\delta$, and define the set

$$W_k = \{w \in S^* \mid w \text{ is freely reduced and } |w| \leq k\}.$$

- Let $p(w)$ denote the path in X associated to the word w .
- For each $w \in W_k$, decide if $p(w)$ is geodesic or not. (Note: W_k is a finite set so this procedure terminates.)
- If $p(w)$ is not geodesic, set $w = u_i$. There exists a word $w' \in W_k$ such that $p(w')$ is geodesic and shares the same end points as $p(w)$. Set $w' = v_i$.
- As a result, $u_i = v_i$, or equivalently, $u_i v_i^{-1} = 1$.
- When the procedure terminates, we obtain a list of relators

$$R = \{u_1 v_1^{-1}, \dots, u_n v_n^{-1}\}.$$

Step 2: Verify the presentation $\langle S \mid R \rangle$ is a Dehn presentation and that $G \cong \langle S \mid R \rangle$.

- If $w =_G 1$, then we want to show that $w \in \ll R \gg$.
- Induct on the length of w .
- Base case: $|w| = 0$, *i.e.*, w is the empty word.
- Inductive step: Suppose all words $w =_G 1$ of length at most L are contained in $\ll R \gg$.
- Let $w =_G 1$ have length $L + 1$. Since $w =_G 1$, $p(w)$ forms a cycle or a closed loop in X .
- By lemma 2, $p(w)$ contains a non-geodesic subarc γ of length less than 8δ .

- By construction, the arc γ corresponds to a word u_i .
- So, $w = w'u_iw'' = w'v_iw''$, but $|v_i| < |u_i|$, so $|w'v_iw''| < L + 1$. By inductive hypothesis, $w'v_iw'' \in \ll R \gg$.
- Now, $w'u_iv_i^{-1}w'^{-1} \in \ll R \gg$ (by definition of normal closure). So,

$$\begin{aligned} w'u_iv_i^{-1}w'^{-1} \cdot w'v_iw'' &= w'u_iv_i^{-1}v_iw'' \\ &= w'u_iw'' \in \ll R \gg . \end{aligned}$$

- Notice that, in the induction, we have also verified that if $w =_G 1$, then w contains a subword of the form u_i , so $\langle S \mid R \rangle$ is indeed a Dehn presentation for G .

□

8 Examples of Hyperbolic Groups

- Any finite group.
- Finitely generated free groups.
- Fundamental group of compact negatively-curved Riemannian manifold.
- Virtually cyclic groups, *e.g.*, infinite dihedral group.⁵
- Virtually free groups, *e.g.*, $F \rtimes H$, where F is free and H is finite, or $H * K$, where H and K are finite.

⁵This is a result of the fact: If H is a finite index subgroup of G , then $G \sim_{QI} H$.

References

- [1] Alexander Yu. Olshanskii. Almost every group is hyperbolic, *J. Algebra Comput.*, 2(1), pp. 1-17, 1992.
- [2] C. Löh, *Geometric group theory* (Universitext). Springer, Cham, 2017, pp. xi+389, An introduction, isbn: 978-3-319-72253-5; 978-3-319-72254-2. doi: 10.1007/978-3-319-72254-2. [Online]. Available: <https://doi-org.uml.idm.oclc.org/10.1007/978-3-319-72254-2>
- [3] G. Hyun, *Hyperbolicity and the word problem*. [Online]. Available: <https://math.uchicago.edu/may/REU2013/REUPapers/Hyun.pdf>
- [4] M. Clay and D. Margalit, Eds., *Office hours with a geometric group theorist*. Princeton University Press, Princeton, NJ, 2017, pp. xii+441, isbn: 978-0-691-15866-2.