

Name: _____

Quiz 7 Solutions

Please write your solutions to the following exercises in the space provided. You should write legibly and fully explain your work.

Good Luck!

(1) Let $A = \begin{bmatrix} 1 & -1 & -1 \\ 0 & 2 & 0 \\ -1 & -1 & 1 \end{bmatrix}$.

- (a) Using the definition of eigenvector, show that $\mathbf{x} = \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix}$ is an eigenvector of A . What is the eigenvalue of \mathbf{x} ? [3 pts]

Solution: The vector \mathbf{x} is non-zero and

$$A\mathbf{x} = \begin{bmatrix} 1 & -1 & -1 \\ 0 & 2 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = 0\mathbf{x}.$$

Since $A\mathbf{x} = 0\mathbf{x}$, \mathbf{x} is an eigenvector of A with eigenvalue $\lambda = 0$.

- (b) $\lambda = 2$ is an eigenvalue of A . Find a basis for the corresponding eigenspace E_2 . What is the geometric multiplicity of the eigenvalue $\lambda = 2$? [8 pts]

Solution: To find a basis for E_2 we find the null space of $(A - 2I)$:

$$[A - 2I \mid \mathbf{0}] = \begin{bmatrix} -1 & -1 & -1 & \mid & 0 \\ 0 & 0 & 0 & \mid & 0 \\ -1 & -1 & -1 & \mid & 0 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 1 & 1 & \mid & 0 \\ 0 & 0 & 0 & \mid & 0 \\ 0 & 0 & 0 & \mid & 0 \end{bmatrix}.$$

Thus,

$$E_2 = \text{span} \left(\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right).$$

A basis for E_2 is

$$\left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

Since $\dim(E_2) = 2$, the geometric multiplicity of $\lambda = 2$ is 2.

- (2) Suppose that $\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = 5$. What is $\begin{vmatrix} a & b & c \\ 2d - 3g & 2e - 3h & 2f - 3i \\ g & h & i \end{vmatrix}$? Be sure to show all of your work. [4 pts]

Solution: Observe that we have the following row reductions:

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow B = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix} \rightarrow C = \begin{bmatrix} a & b & c \\ 2d - 3g & 2e - 3h & 2f - 3i \\ g & h & i \end{bmatrix}.$$

By Theorem 4.3,

$$\det(C) = \det(B) = 2 \det(A) = 2(5) = 10.$$

- (3) Suppose that A and B are $n \times n$ matrices with $\det(A) = 2$ and $\det(B) = -5$. Find $\det(B^{-1}A)$. Be sure to show all of your work. [5 pts]

Solution:

$$\det(B^{-1}A) = \det(B^{-1}) \det(A) = \frac{1}{\det(B)} \det(A) = -\frac{2}{5}.$$